

Too good to be true? Spotting fake trials in the literature

Jack Wilkinson, Centre for Biostatistics, University of Manchester.



@jdwilko.bsky.social

Some of the research discussed in this presentation is funded by the NIHR Research for Patient Benefit programme (NIHR203568, NIHR303741). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

FUNDED BY

Disclaimer

I am not accusing anyone of fraud or other forms of misconduct. A study might be 'problematic' or 'untrustworthy' due to honest errors.

Declaration of interest

I am Stats Editor for BJOG, Cochrane Gynaecology and Fertility, Fertility and Sterility, Journal of Hypertension. I undertake integrity investigations for various journals, and sometimes I am paid for this.

Systematic reviews of health interventions

Cochrane Database of Systematic Reviews | Review - Intervention


Probiotic treatment for women with gestational diabetes to improve maternal and infant health and well-being

✉ Karaponi AM Okesene-Gafa, Abigail E Moore, Vanessa Jordan, Lesley McCowan, Caroline A Crowther

Authors' declarations of interest

Version published: 24 June 2020 Version history

<https://doi.org/10.1002/14651858.CD012970.pub2>



	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): All outcomes	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
Ahmadi 2016	+	?	+	+	+	?	+
Badehnoosh 2018	+	?	+	+	+	?	+
Hajifaraji 2017	+	?	+	+	+	?	+
Jafarnejad 2016	+	?	+	+	+	?	+
Karamali 2016	+	?	+	+	+	?	+
Karamali 2018	+	?	+	+	+	?	+
Kijmanawat 2019	+	?	+	+	+	?	+
Lindsay 2015	+	+	+	+	+	?	+
Nabhani 2018	+	?	+	+	+	?	+

Systematic reviews of health interventions

Cochrane Database of Systematic Reviews | Review - Intervention


Probiotic treatment for women with gestational diabetes to improve maternal and infant health and well-being

✉ Karaponi AM Okesene-Gafa, Abigail E Moore, Vanessa Jordan, Lesley McCowan, Caroline A Crowther

Authors' declarations of interest

Version published: 24 June 2020 Version history

<https://doi.org/10.1002/14651858.CD012970.pub2>



	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): All outcomes	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
Ahmadi 2016	+	?	+	+	+	?	+
Badehnoosh 2018	+	?	+	+	+	?	+
Hajifaraji 2017	+	?	+	+	+	?	+
Jafarnejad 2016	+	?	+	+	+	?	+
Karamali 2016	+	?	+	+	+	?	+
Karamali 2018	+	?	+	+	+	?	+
Kijmanawat 2019	+	?	+	+	+	?	+
Lindsay 2015	+	+	+	+	+	?	+
Nabhani 2018	+	?	+	+	+	?	+

Systematic reviews of health interventions

Cochrane Database of Systematic Reviews | Review - Intervention

Probiotic treatment for women with gestational diabetes to improve maternal and infant health and well-being

✉ Karaponi AM Okesene-Gafa, Abigail E Moore, Vanessa Jordan, Lesley McCowan, Caroline A Crowther

Authors' declarations of interest

Version published: 24 June 2020 Version history

<https://doi.org/10.1002/14651858.CD012970.pub2>


Study or Subgroup	Probiotics		Placebo		Weight	Risk Ratio M-H, Fixed, 95% CI
	Events	Total	Events	Total		
Badehnoosh 2018	2	30	2	30	25.0%	1.00 [0.15 , 6.64]
Karamali 2018	4	30	4	30	50.0%	1.00 [0.28 , 3.63]
Lindsay 2015	6	68	2	68	25.0%	3.00 [0.63 , 14.34]
Total (95% CI)		128		128	100.0%	1.50 [0.64 , 3.53]

Total events: 12 8

Heterogeneity: $\text{Chi}^2 = 1.31$, $\text{df} = 2$ ($P = 0.52$); $I^2 = 0\%$

Test for overall effect: $Z = 0.93$ ($P = 0.35$)

Test for subgroup differences: Not applicable



	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): All outcomes	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
Ahmadi 2016	+	?	+	+	+	?	+
Badehnoosh 2018	+	?	+	+	+	?	+
Hajifaraji 2017	+	?	+	+	+	?	+
Jafarnejad 2016	+	?	+	+	+	?	+
Karamali 2016	+	?	+	+	+	?	+
Karamali 2018	+	?	+	+	+	?	+
Kijmanawat 2019	+	?	+	+	+	?	+
Lindsay 2015	+	+	+	+	+	?	+
Nabhani 2018	+	?	+	+	+	?	+

Vitamin K and the Prevention of Fractures

Systematic Review and Meta-analysis of Randomized Controlled Trials

*Sarah Cockayne, MSc; Joy Adamson, PhD; Susan Lanham-New, PhD; Martin J. Shearer, PhD, MRCPath;
Simon Gilbody, DPhil; David J. Torgerson, PhD*

Ivermectin for Prevention and Treatment of COVID-19 Infection: A Systematic Review, Meta- analysis, and Trial Sequential Analysis to Inform Clinical Guidelines

 Bryant, Andrew MSc^{1*}; Lawrie, Theresa A. MBBCh, PhD²; Dowswell, Therese PhD²; Fordham, Edmund J. PhD²; Mitchell, Scott MBChB, MRCS³; Hill, Sarah R. PhD¹; Tham, Tony C. MD, FRCP⁴

Psychological therapies for the management of chronic pain (excluding headache) in adults (Review)

Williams ACDC, Fisher E, Hearn L, Eccleston C

Vitamin K and the Prevention of Fractures

Systematic Review and Meta-analysis of Randomized Controlled Trials

Sarah Cockayne, MSc; Joy Adamson, PhD; Susan Lanham-New, PhD; Martin J. Shearer, PhD, MRCPath; Simon Gilbody, DPhil; David J. Torgerson, PhD

Ivermectin for Prevention and Treatment of COVID-19 Infection: A Systematic Review, Meta-analysis, and Trial Sequential Analysis to Inform Clinical Guidelines

 Bryant, Andrew MSc^{1*}; Lawrie, Theresa A. MBBCh, PhD²; Dowswell, Therese PhD²; Fordham, Edmund J. PhD²; Mitchell, Scott MBChB, MRCS³; Hill, Sarah R. PhD¹; Tham, Tony C. MD, FRCP⁴

Psychological therapies for the management of chronic pain (excluding headache) in adults (Review)

Williams ACDC, Fisher E, Hearn L, Eccleston C

3 out of 5 trials subsequently identified as **fake**.

Suggested impressive benefit on mortality due to **fraudulent** trials.

3 trials from one investigator suggested to be **implausible** (huge effects, no attrition), 2 **retracted**.

Systematic reviews: Fake data to patient care pipeline

1

**Attempt to identify all RCTs
on the review topic**

- Problematic trials will be included



2

**Critically appraise study
methodology, include in
meta-analysis**

- Assess risk of bias
- But do not consider authenticity
- Many (not all) fake trials report sound methods



3

**Make conclusions,
recommendations, on
basis of evidence**

- SRs seen as gold standard
- Included in guidelines
- Influence patient care



How big is the problem?

Difficult to estimate!

[Carlisle 2020](#): 26% RCTs submitted to Anaesthesia “critically flawed by false data”.

[Gopalakrishna 2022](#): 10% of researchers in life/medical sciences have committed fabrication or falsification.

[Xu 2025](#): Identified 3902 meta-analyses in 847 systematic reviews synthesising retracted trials.

EDITORIAL

When beauty is but skin deep: dealing with problematic studies in systematic reviews

Stephanie L Boughton, Jack Wilkinson, Lisa Bero

Managing potentially problematic studies

<https://bit.ly/3SsJO9F>

EDITORIAL

When beauty is but skin deep: dealing with problematic studies in systematic reviews

Stephanie L Boughton, Jack Wilkinson, Lisa Bero

Managing potentially problematic studies

<https://bit.ly/3SsJO9F>

- Do not include problematic studies in the systematic review.

EDITORIAL

When beauty is but skin deep: dealing with problematic studies in systematic reviews


Stephanie L Boughton, Jack Wilkinson, Lisa Bero

Managing potentially problematic studies

<https://bit.ly/3SsJO9F>

- Do not include problematic studies in the systematic review.
- But how can we identify problematic studies in the first place?

Can't we just use retraction status?



It typically takes a long time for problematic studies to be retracted


Many are never retracted

Example (left): outcome 5 years after reporting 172 problematic clinical trials from one research group

So we cannot rely on retraction status alone as an indicator of trustworthiness

Do we need a tool to identify fraud and misconduct?

Our view: it is neither necessary nor desirable to develop a “diagnostic test for fraud”



```
graph TD; A[Our view: it is neither necessary nor desirable to develop a “diagnostic test for fraud”] --> B[Legal consequences, harms]; B --> C[What matters it the trustworthiness of a study (example)]; C --> D[A trustworthiness tool should help the user make a judgement about a study’s trustworthiness, and to articulate a basis for that judgement]; D --> E[Wilkinson and Tovey 2025, JCE: doi.org/10.1016/j.jclinepi.2025.111670];
```

Legal consequences, harms

What matters it the trustworthiness of a study (example)

A trustworthiness tool should help the user make a judgement about a study’s trustworthiness, and to articulate a basis for that judgement

Wilkinson and Tovey 2025, JCE:
doi.org/10.1016/j.jclinepi.2025.111670

Table I. Patients' characteristics.

	Group A (minimal stimulation) (<i>n</i> = 58)	Group B (CC) (<i>n</i> = 55)	<i>p</i> Value
Age (years)	27.7 ± 2.4	26.8 ± 2.3	0.57
Duration of infertility (years)	1.88 ± 0.7	2.14 ± 0.8	0.71
BMI (kg/m ²)	24.9 ± 2.3	25.3 ± 2.6	0.62
Waist-to-hip ratio	0.8 ± 0.04	0.8 ± 0.05	0.38
Menstrual cycle			
Oligomenorrhea	54 (93.1%)	51 (92.7%)	0.43
Amenorrhea	4 (6.9%)	4 (7.3%)	0.61
Hyperandrogenism	26 (44.8%)	23 (41.8%)	0.74
LH (mIU/mL)	12.9 ± 2.3	13.4 ± 2.4	0.52
FSH (mIU/mL)	5.6 ± 1.1	5.7 ± 1.3	0.72
LH/FSH ratio	2.6 ± 1.1	2.6 ± 1.2	0.42
Testosterone (ng/mL)	0.94 ± 0.21	0.92 ± 0.24	0.77
SHBG (nmol/L)	33.62 ± 3.6	35.53 ± 3.1	0.48
Fasting glucose (mg/dL)	91.6 ± 1.5	89.7 ± 1.4	0.76
Fasting insulin (μU/mL)	10.7 ± 2.7	11.2 ± 3.1	0.57
Fasting glucose/insulin ratio	7.7 ± 3.7	7.2 ± 3.6	0.33
Ovarian volume (mL)	11.3 ± 2.5	11.4 ± 2.4	0.22

BMI, Body mass index; CC, clomiphene citrate; FSH, follicle-stimulating hormone;


Table I. Patients' characteristics.

	Group A (minimal stimulation) (<i>n</i> = 58)	Group B (CC) (<i>n</i> = 55)	<i>p</i> Value
Age (years)	27.7 ± 2.4	26.8 ± 2.3	0.57
Duration of infertility (years)	1.88 ± 0.7	2.14 ± 0.8	0.71
BMI (kg/m ²)	24.9 ± 2.3	25.3 ± 2.6	0.62
Waist-to-hip ratio	0.8 ± 0.04	0.8 ± 0.05	0.38
Menstrual cycle			
Oligomenorrhea	54 (93.1%)	51 (92.7%)	0.43
Amenorrhea	4 (6.9%)	4 (7.3%)	0.61
Hyperandrogenism	26 (44.8%)	23 (41.8%)	0.74
LH (mIU/mL)	12.9 ± 2.3	13.4 ± 2.4	0.52
FSH (mIU/mL)	5.6 ± 1.1	5.7 ± 1.3	0.72
LH/FSH ratio	2.6 ± 1.1	2.6 ± 1.2	0.42
Testosterone (ng/mL)	0.94 ± 0.21	0.92 ± 0.24	0.77
SHBG (nmol/L)	33.62 ± 3.6	35.53 ± 3.1	0.48
Fasting glucose (mg/dL)	91.6 ± 1.5	89.7 ± 1.4	0.76
Fasting insulin (μU/mL)	10.7 ± 2.7	11.2 ± 3.1	0.57
Fasting glucose/insulin ratio	7.7 ± 3.7	7.2 ± 3.6	0.33
Ovarian volume (mL)	11.3 ± 2.5	11.4 ± 2.4	0.22

BMI, Body mass index; CC, clomiphene citrate; FSH, follicle-stimulating hormone;

Do we need a tool to identify fraud and misconduct?

Our view: it is neither necessary nor desirable to develop a “diagnostic test for fraud”



```
graph TD; A[Our view: it is neither necessary nor desirable to develop a “diagnostic test for fraud”] --> B[Legal consequences, harms]; B --> C[What matters it the trustworthiness of a study (example)]; C --> D[A trustworthiness tool should help the user make a judgement about a study’s trustworthiness, and to articulate a basis for that judgement]; D --> E[Wilkinson and Tovey 2025, JCE: doi.org/10.1016/j.jclinepi.2025.111670];
```

Legal consequences, harms

What matters it the trustworthiness of a study (example)

A trustworthiness tool should help the user make a judgement about a study’s trustworthiness, and to articulate a basis for that judgement

Wilkinson and Tovey 2025, JCE:
doi.org/10.1016/j.jclinepi.2025.111670



INveStigating ProblEmatic Clinical Trials in Systematic Reviews

Aim: To develop a tool for identifying problematic randomised controlled trials in the context of health systematic reviews.



INveStigating ProBlEmatic Clinical Trials in Systematic Reviews

Aim: To develop a tool for identifying problematic randomised controlled trials in the context of health systematic reviews.

Stage 1: Assemble list
of checks for
problematic studies
(previous studies, new
survey of 71 experts):
76 checks identified



INveStigating ProblEmatic Clinical Trials in Systematic Reviews

Aim: To develop a tool for identifying problematic randomised controlled trials in the context of health systematic reviews.

Stage 1: Assemble list
of checks for
problematic studies
(previous studies, new
survey of 71 experts):
76 checks identified



Stage 2: Apply list of
checks to 95 RCTs in
50 Cochrane Reviews
(feasibility, impact)

INveStigating ProbleMatic Clinical Trials in Systematic Reviews

Aim: To develop a tool for identifying problematic randomised controlled trials in the context of health systematic reviews.


Stage 1: Assemble list
of checks for
problematic studies
(previous studies, new
survey of 71 experts):
76 checks identified

Stage 2: Apply list of
checks to 95 RCTs in
50 Cochrane Reviews
(feasibility, impact)

Stage 3: Delphi survey
of 158 experts and
users (**26 checks** are
backed by broad
consensus)


INveStigating ProbleMatic Clinical Trials in Systematic Reviews


Aim: To develop a tool for identifying problematic randomised controlled trials in the context of health systematic reviews.





INveStigating ProblEmatic Clinical Trials in Systematic Reviews

Aim: To develop a tool for identifying problematic randomised controlled trials in the context of health systematic reviews.









Inspecting post-publication notices

3 checks

Overall domain judgement:

No concerns, some concerns,
serious concerns

Inspecting conduct, governance and transparency

5 checks

Overall domain judgement:

No concerns, some concerns,
serious concerns

Inspecting text and figures

2 checks

Overall domain judgement:

No concerns, some concerns,
serious concerns

Inspecting results in the study


11 checks

Overall domain judgement:

No concerns, some concerns,
serious concerns

Overall study judgement:

No concerns, some concerns,
serious concerns



Inspecting post-publication notices
3 checks



Overall domain judgement:
No concerns, some concerns,
serious concerns

Inspecting conduct, governance and transparency
5 checks



Overall domain judgement:
No concerns, some concerns,
serious concerns

Inspecting text and figures
2 checks




Overall domain judgement:
No concerns, some concerns,
serious concerns

Inspecting results in the study
11 checks



Overall domain judgement:
No concerns, some concerns,
serious concerns



Overall study judgement:
No concerns, some concerns,
serious concerns

Probiotic treatment for women with gestational diabetes to improve maternal and infant health and well-being

✉ Karaponi AM Okesene-Gafa, Abigail E Moore, Vanessa Jordan, Lesley McCowan, Caroline A Crowther

Authors' declarations of interest

Version published: 24 June 2020 Version history

<https://doi.org/10.1002/14651858.CD012970.pub2>

Study or Subgroup	Probiotics		Placebo		Weight	Risk Ratio	Risk Ratio
	Events	Total	Events	Total		M-H, Fixed, 95% CI	M-H, Fixed, 95% CI
Badehnoosh 2018	2	30	2	30	25.0%	1.00 [0.15 , 6.64]	
Karamali 2018	4	30	4	30	50.0%	1.00 [0.28 , 3.63]	
Lindsay 2015	6	68	2	68	25.0%	3.00 [0.63 , 14.34]	
Total (95% CI)		128		128	100.0%	1.50 [0.64 , 3.53]	
Total events:	12		8				
Heterogeneity: Chi ² = 1.31, df = 2 (P = 0.52); I ² = 0%							
Test for overall effect: Z = 0.93 (P = 0.35)							
Test for subgroup differences: Not applicable							

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): All outcomes	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
Ahmadi 2016	+	?	+	+	+	?	+
Badehnoosh 2018	+	?	+	+	+	?	+
Hajifaraji 2017	+	?	+	+	+	?	+
Jafarnejad 2016	+	?	+	+	+	?	+
Karamali 2016	+	?	+	+	+	?	+
Karamali 2018	+	?	+	+	+	?	+
Kijmanawat 2019	+	?	+	+	+	?	+
Lindsay 2015	+	+	+	+	+	?	+
Nabhani 2018	+	?	+	+	+	?	+

Probiotic treatment for women with gestational diabetes to improve maternal and infant health and well-being

✉ Karaponi AM Okesene-Gafa, Abigail E Moore, Vanessa Jordan, Lesley McCowan, Caroline A Crowther

Authors' declarations of interest

Version published: 24 June 2020 Version history

<https://doi.org/10.1002/14651858.CD012970.pub2>

Study or Subgroup	Probiotics		Placebo		Weight	Risk Ratio	Risk Ratio
	Events	Total	Events	Total		M-H, Fixed, 95% CI	M-H, Fixed, 95% CI
Badehnoosh 2018	2	30	2	30	25.0%	1.00 [0.15 , 6.64]	
Karamali 2018	4	30	4	30	50.0%	1.00 [0.28 , 3.63]	
Lindsay 2015	6	68	2	68	25.0%	3.00 [0.63 , 14.34]	
Total (95% CI)		128		128	100.0%	1.50 [0.64 , 3.53]	
Total events:	12		8				
Heterogeneity: Chi ² = 1.31, df = 2 (P = 0.52); I ² = 0%							
Test for overall effect: Z = 0.93 (P = 0.35)							
Test for subgroup differences: Not applicable							

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): All outcomes	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
Ahmadi 2016	+	?	+	+	+	?	+
Badehnoosh 2018	+	?	+	+	+	?	+
Hajifaraji 2017	+	?	+	+	+	?	+
Jafarnejad 2016	+	?	+	+	+	?	+
Karamali 2016	+	?	+	+	+	?	+
Karamali 2018	+	?	+	+	+	?	+
Kijmanawat 2019	+	?	+	+	+	?	+
Lindsay 2015	+	+	+	+	+	?	+
Nabhani 2018	+	?	+	+	+	?	+


Probiotic treatment for women with gestational diabetes to improve maternal and infant health and well-being

✉ Karaponi AM Okesene-Gafa, Abigail E Moore, Vanessa Jordan, Lesley McCowan, Caroline A Crowther

Authors' declarations of interest

Version published: 24 June 2020 Version history

<https://doi.org/10.1002/14651858.CD012970.pub2>



Random sequence generation (selection bias)
Allocation concealment (selection bias)
Blinding of participants and personnel (performance bias): All outcomes
Blinding of outcome assessment (detection bias): All outcomes
Incomplete outcome data (attrition bias): All outcomes
Selective reporting (reporting bias)
Other bias

Ahmadi 2016	+	?	+	+	+	?	+
Badehnoosh 2018	+	?	+	+	+	?	+
Hajifaraji 2017	+	?	+	+	+	?	+
Jafarnejad 2016	+	?	+	+	+	?	+
Karamali 2016	+	?	+	+	+	?	+
Karamali 2018	+	?	+	+	+	?	+
Kijmanawat 2019	+	?	+	+	+	?	+
Lindsay 2015	+	+	+	+	+	?	+
Nabhani 2018	+	?	+	+	+	?	+

Inspecting post-publication notices

Does the study have an associated retraction?

This study has recently been **retracted**.

RETRACTED ARTICLE: The effects of probiotic supplementation on biomarkers of inflammation, oxidative stress and pregnancy outcomes in gestational diabetes

Both the journal editorial team and the external statistical reviewer were unable to confirm the integrity of the trial design and the main outcome of the external review was that the article's conclusions are unreliable. Therefore, as the editorial team no longer have confidence in the reported conclusions the decision has been made to retract the article.

Does consideration of other studies from members of the research team highlight causes for concern?

Search on last author on Retraction Watch database: <http://retractiondatabase.org/>

Author(s):	<input type="text" value="Asemi, Zatollah"/>	Country(s):	<input type="text"/>
Title:	<input type="text" value="Type to search"/>		
Reason(s) for Retraction:	<input type="text"/>		
Subject(s):	<input type="text"/>	Article Type(s):	<input type="text"/>
Journal:	<input type="text"/>		
Publisher:	<input type="text"/>		
Affiliation(s):	<input type="text"/>		
Notes:	<input type="text"/>		
URL:	<input type="text"/>		

Does consideration of other studies from members of the research team highlight causes for concern?

Search on last author on Retraction Watch database: <http://retractiondatabase.org/>

Author(s):	<input type="text" value="Asemi, Zatollah"/>	Country(s):	<input type="text"/>
Title:	<input type="text" value="Type to search"/>		
Reason(s) for Retraction:	<input type="text"/>		
Subject(s):	<input type="text"/>	Article Type(s):	<input type="text"/>
Journal:	<input type="text"/>		
Publisher:	<input type="text"/>		
Affiliation(s):	<input type="text"/>		
Notes:	<input type="text"/>		
URL:	<input type="text"/>		

Retraction or Other Notices			
Title/Subject(s)/Journal --- Publisher/Affiliation(s)/Retraction Watch Post URL(s)			
50 Items Displayed Out of 72 Item(s) Found			

Inspecting conduct, governance and transparency

Are there important inconsistencies between the publication and the registration documents?

- Retrospective registration, with **various inconsistencies** (e.g. recruitment period 6 months vs 1 month)
- Description of the control arm differs between paper and registration (later changed to match):

Description - English

Control group: Placebo (Barij Essence, Kashan, Iran), daily, for 6 weeks orally.

Control group: Placebo (Tak Gen Zist, Tehran, Iran), daily, for 6 weeks orally.







Description - Persian

گروه کنترل: پلاسبو (باریج اسانس، کاشان، ایران)، روزانه، به صورت خوراکی برای 6 هفته

گروه کنترل: پلاسبو (تک ژن زیست، تهران، ایران)، روزانه، به صورت خوراکی برای 6 هفته

Inspecting text and figures

Is there evidence of manipulation or duplication of figures?



 Active Laser Group
  Placebo Laser Group

Inspecting results in the study

Are the means and variances of integer data impossible?

1-min Apgar score	8.93 ± 0.25	8.96 ± 0.18	.561
5-min Apgar score	9.93 ± 0.18	9.96 ± 0.18	.561
Newborns' hyperbilirubinemia (%)	8 (26.7)	2 (6.7)	.080 ^b
Newborns' hospitalization (%)	8 (26.7)	2 (6.7)	.080 ^b
Newborns' hypoglycemia (%)	3 (10.0)	2 (6.7)	>.999 ^b


Inspecting results in the study

Are the means and variances of integer data impossible?

1-min Apgar score	8.93 ± 0.25	8.96 ± 0.18	.561
5-min Apgar score	9.93 ± 0.18	9.96 ± 0.18	.561
Newborns' hyperbilirubinemia (%)	8 (26.7)	2 (6.7)	.080 ^b
Newborns' hospitalization (%)	8 (26.7)	2 (6.7)	.080 ^b
Newborns' hypoglycemia (%)	3 (10.0)	2 (6.7)	>.999 ^b

- Apgar score is a variable which only takes **integer values** (1,2,3,4,5,6,7,8,9,10).
- The highlighted values **cannot occur** for the group sizes in the study.
- 'GRIM' and 'GRIMMER'


Hypertensive disorders



Are any outcome data, including estimated treatment effects, implausible?

- Karamali 2018 is another trial from same group.

Caesarian deliveries



- Almost identical risk ratios in the two studies (2 here, several other outcomes have identical effect estimates).

Domain	Check	Result
Inspecting post-publication notices	<i>Does the study have an associated retraction?</i>	Yes , this article has been retracted
	<i>Do other studies by the research team highlight causes for concern?</i>	Yes , the senior author has more than 50 notices on Retraction Watch
Inspecting conduct, governance, and transparency	<i>Are there important inconsistencies between the publication and the registration documents?</i>	Yes , various discrepancies, including recruitment dates and description of control intervention
Inspecting text and figures	<i>Is there evidence of manipulation or duplication of figures?</i>	NA , no figures in the manuscript
Inspecting results in the study	<i>Are the means and variances of integer data impossible?</i>	Yes , there are impossible means and variances for Apgar scores
	<i>Are any outcome data, including estimated treatment effects, implausible?</i>	Yes , results essentially identical in another RCT from this team, across multiple measures

Applying the draft INSPECT-SR tool would lead to **serious concerns** about the study's trustworthiness.

🔔 Follow this preprint

INSPECT-SR: a tool for assessing trustworthiness of randomised controlled trials

 Jack Wilkinson, Calvin Heal, Ella Flemmyng, Georgios A. Antoniou, Tony Aburrow, Zarko Alfirevic, Alison Avenell, Virginia Barbour, Vincenzo Berghella, Dorothy V. M. Bishop, Esmée M Bordewijk, Nicholas J. L. Brown, Jana Christopher, Mike Clarke,  Darren Dahly, Jane Dennis, Patrick Dicker, Jo Dumville, Helen Frankish, Andrew Grey, Steph Grohmann, Lyle C. Gurrin, Jill A. Hayden, James A.J. Heathers,  Kylie E Hunter,  Ian Hussey,  Lukas Jung, Emily Lam, Toby J. Lasserson, Sarah Lensen, Tianjing Li, Wentao Li, Jianping Liu, Elizabeth Loder, Andreas Lundh, Gideon Meyerowitz-Katz, Ben W. Mol,  Florian Naudet, Anna Noel-Storr,  Neil E. O'Connell, Lisa Parker, Rita F. Redberg, Barbara K. Redman, Rachel Richardson, Anna Lene Seidler, Kyle Sheldrick, Emma Sydenham, Madelon van Wely, Colby J. Vorland, Rui Wang, Stephanie Weibel, Matthias Wjst, Lisa Bero,  Jamie J. Kirkham

doi: <https://doi.org/10.1101/2025.09.03.25334905>

Upcoming online training workshop:

- **July 23rd** 10am-12pm (London, UK)

Booking link:

<https://www.trybooking.com/uk/GNCQ>

- INSPECT-SR is now available
- Endorsed by Cochrane:
<https://www.cochrane.org/authors/methods-cochrane>
- An editable template and detailed guidance document are provided: <https://osf.io/b74wj> and inspect.sr



***Retracted:* Randomized controlled trial of elevation of the fetal head with a fetal pillow during cesarean delivery at full cervical dilatation**

 **Correction(s) for this article** ▾

 **Retraction(s) for this article** ▾

Subrata L. Seal, Alok Dey, Sannyashi C. Barman, Gourisankar Kamilya, Joydev Mukherji,
Joseph L. Onwude✉

- RCT retracted – National Institute for Health and Care Excellence (NICE) subsequently reversed recommendation of this device.
- Royal College of Obstetrics and Gynaecology retracted Scientific Impact Paper
- See <https://doi.org/10.1111/1471-0528.18004>

- INSPECT-SR would have flagged problems ahead of retraction
- E.g. Numerical anomalies in the abstract and results

Results

Overall, 120 women were assigned to each group. Major uterine wound extensions occurred in 6 (5.0%) women in the Fetal Pillow group and 39 (32.5%) in the control group (relative risk 0.23, 95% confidence interval 0.11–0.48).

***Retracted:* Randomized controlled trial of elevation of the fetal head with a fetal pillow during cesarean delivery at full cervical dilatation**

! [Correction\(s\) for this article](#) ▾

! [Retraction\(s\) for this article](#) ▾

Subrata L. Seal, Alok Dey, Sannyashi C. Barman, Gourisankar Kamilya, Joydev Mukherji,
Joseph L. Onwude✉

- RCT retracted – National Institute for Health and Care Excellence (NICE) subsequently reversed recommendation of this device.
- Royal College of Obstetrics and Gynaecology retracted Scientific Impact Paper
- See <https://doi.org/10.1111/1471-0528.18004>

- INSPECT-SR would have flagged problems ahead of retraction
- E.g. Numerical anomalies in the abstract and results

Results

Overall, 120 women were assigned to each group. Major uterine wound extensions occurred in 6 (5.0%) women in the Fetal Pillow group and 39 (32.5%) in the control group (relative risk 0.23, 95% confidence interval 0.11–0.48).

Are there errors in statistical results?

Table I. Patients' characteristics.

	Group A (minimal stimulation) (<i>n</i> = 58)	Group B (CC) (<i>n</i> = 55)	<i>p</i> Value
Age (years)	27.7 ± 2.4	26.8 ± 2.3	0.57
Duration of infertility(years)	1.88 ± 0.7	2.14 ± 0.8	0.71
BMI (kg/m ²)	24.9 ± 2.3	25.3 ± 2.6	0.62
Waist-to-hip ratio	0.8 ± 0.04	0.8 ± 0.05	0.38
Menstrual cycle			
Oligomenorrhea	54 (93.1%)	51 (92.7%)	0.43
Amenorrhea	4 (6.9%)	4 (7.3%)	0.61
Hyperandrogenism	26 (44.8%)	23 (41.8%)	0.74
LH (mIU/mL)	12.9 ± 2.3	13.4 ± 2.4	0.52
FSH (mIU/mL)	5.6 ± 1.1	5.7 ± 1.3	0.72
LH/FSH ratio	2.6 ± 1.1	2.6 ± 1.2	0.42
Testosterone (ng/mL)	0.94 ± 0.21	0.92 ± 0.24	0.77
SHBG (nmol/L)	33.62 ± 3.6	35.53 ± 3.1	0.48
Fasting glucose (mg/dL)	91.6 ± 1.5	89.7 ± 1.4	0.76
Fasting insulin (μU/mL)	10.7 ± 2.7	11.2 ± 3.1	0.57
Fasting glucose/insulin ratio	7.7 ± 3.7	7.2 ± 3.6	0.33
Ovarian volume (mL)	11.3 ± 2.5	11.4 ± 2.4	0.22

BMI, Body mass index; CC, clomiphene citrate; FSH, follicle-stimulating hormone;

- Look at waist-to-hip ratio, which has been analysed by t-test.
- Do you think there are any problems with these results?



Are there errors in statistical results?

Table I. Patients' characteristics.

	Group A (minimal stimulation) (<i>n</i> = 58)	Group B (CC) (<i>n</i> = 55)	<i>p</i> Value
Age (years)	27.7 ± 2.4	26.8 ± 2.3	0.57
Duration of infertility (years)	1.88 ± 0.7	2.14 ± 0.8	0.71
BMI (kg/m ²)	24.9 ± 2.3	25.3 ± 2.6	0.62
Waist-to-hip ratio	0.8 ± 0.04	0.8 ± 0.05	0.38
Menstrual cycle			
Oligomenorrhea	54 (93.1%)	51 (92.7%)	0.43
Amenorrhea	4 (6.9%)	4 (7.3%)	0.61
Hyperandrogenism	26 (44.8%)	23 (41.8%)	0.74
LH (mIU/mL)	12.9 ± 2.3	13.4 ± 2.4	0.52
FSH (mIU/mL)	5.6 ± 1.1	5.7 ± 1.3	0.72
LH/FSH ratio	2.6 ± 1.1	2.6 ± 1.2	0.42
Testosterone (ng/mL)	0.94 ± 0.21	0.92 ± 0.24	0.77
SHBG (nmol/L)	33.62 ± 3.6	35.53 ± 3.1	0.48
Fasting glucose (mg/dL)	91.6 ± 1.5	89.7 ± 1.4	0.76
Fasting insulin (μU/mL)	10.7 ± 2.7	11.2 ± 3.1	0.57
Fasting glucose/insulin ratio	7.7 ± 3.7	7.2 ± 3.6	0.33
Ovarian volume (mL)	11.3 ± 2.5	11.4 ± 2.4	0.22

BMI, Body mass index; CC, clomiphene citrate; FSH, follicle-stimulating hormone;



- Look at waist-to-hip ratio, which has been analysed by t-test.
- Do you think there are any problems with these results?
- We *might* expect a p-value of 1 because the means are equal. However!
- Must remember that the summary stats in the paper are **rounded**.
- These data are consistent with very small p-values (e.g. $p < 0.00001$) so there is no contradiction here.
- Could have means of 0.75 and 0.85, smaller SDs.
- Have to check whether p-values are **consistent with** rounded summary data (see INSPECT-SR guidance check 4.9)

Are there errors in statistical results?

Table I. Patients' characteristics.

	Group A (minimal stimulation) (n = 58)	Group B (CC) (n = 55)	p Value
Age (years)	27.7 ± 2.4	26.8 ± 2.3	0.57
Duration of infertility (years)	1.88 ± 0.7	2.14 ± 0.8	0.71
BMI (kg/m ²)	24.9 ± 2.3	25.3 ± 2.6	0.62
Waist-to-hip ratio	0.8 ± 0.04	0.8 ± 0.05	0.38
Menstrual cycle			
Oligomenorrhea	54 (93.1%)	51 (92.7%)	0.43
Amenorrhea	4 (6.9%)	4 (7.3%)	0.61
Hyperandrogenism	26 (44.8%)	23 (41.8%)	0.74
LH (mIU/mL)	12.9 ± 2.3	13.4 ± 2.4	0.52
FSH (mIU/mL)	5.6 ± 1.1	5.7 ± 1.3	0.72
LH/FSH ratio	2.6 ± 1.1	2.6 ± 1.2	0.42
Testosterone (ng/mL)	0.94 ± 0.21	0.92 ± 0.24	0.77
SHBG (nmol/L)	33.62 ± 3.6	35.53 ± 3.1	0.48
Fasting glucose (mg/dL)	91.6 ± 1.5	89.7 ± 1.4	0.76
Fasting insulin (μU/mL)	10.7 ± 2.7	11.2 ± 3.1	0.57
Fasting glucose/insulin ratio	7.7 ± 3.7	7.2 ± 3.6	0.33
Ovarian volume (mL)	11.3 ± 2.5	11.4 ± 2.4	0.22

BMI, Body mass index; CC, clomiphene citrate; FSH, follicle-stimulating hormone;

It turns out that loads of the results in the paper **are incompatible** with the p-values even taking rounding into account.

All of the results in red are not possible given these summary statistics.

Similar problems in outcome table, including primary outcome:

Clinical pregnancy per patient p = 0.77 vs 0.36

Clinical pregnancy per cycle p = 0.76 vs 0.23

Being able to exactly reproduce p-values from rounded summary data might actually indicate a problem

Inflammatory variables	Group A: control (n = 30)	Group B: esmolol (n = 30)	t test/ χ^2 *	P value
MAP mmHg	70.2 ± 6.42	72.3 ± 6.7	1.240	0.2201
HR rate/min	115.75 ± 12.86	113.4 ± 12.78	− 0.710	0.4806
CVP CmH2o	11.7 ± 4.1	11.6 ± 2.6	− 0.113	0.9106
ScvO2 %	76.2 ± 5.92	74.3 ± 4.8	− 1.365	0.1774
Lactate	5.3 ± 1.35	5.2 ± 1.6	− 0.262	0.7945
APACHE II	23.5 ± 6.2	24.2 ± 5.4	0.466	0.6427
SOFA score	9.2 ± 3.2	8.8 ± 3.5	− 0.462	0.6458

Data presented as mean ± SD

P value > 0.05 NS; *P value < 0.05 S; **P value < 0.001 HS

There was no statistically significant difference between both groups according to admission data as shown in Table 2

- Given that summary data are rounded, we don't expect to be able to reproduce p-values exactly.
- If we *can* reproduce them all exactly, that might suggest no underlying data were analysed to produce the results
- That could suggest that there is no underlying data. Alternative explanation: they calculated summary statistics, and then fed those into a calculator (an odd thing to do)
- Example: the p-values can be reproduced to 4dp from the rounded summary data

	A	B	C	D	E	F
1	Surname and Name	Sex	Birthdate	Eye	Date of Surgery	Age at Surgery
2	Alison Green	M	05/06/1976	RE	11/08/2008	32
3	Jared Friedman	F	13/04/1997	LE	13/04/2015	17
4	Jessica Cohen	M	21/03/1976	RE	28/12/2008	32
5	Sara Young DDS	F	12/05/1964	LE	03/04/2010	45
6	Whitney Douglas	M	26/01/1974	RE	02/01/2009	34
7	Daniel Watkins	M	13/11/1987	RE	03/11/2006	18
8	Jennifer Johnson	M	05/08/1952	RE	31/07/2010	57
9	Tara Thompson	M	16/06/1999	RE	16/06/2017	18
10	Jennifer Dixon	M	30/11/1977	RE	14/03/2007	29
11	Susan Howe	M	19/04/1955	LE	04/04/2005	49
12	Daniel McCarthy	F	11/10/1989	LE	30/04/2009	19
13	Bianca Boyd	M	07/11/1951	RE	02/05/2005	53
14	Lisa Kaufman	M	23/11/1996	RE	23/11/2014	17
15	Michael Stephen	M	06/10/1972	RE	21/07/2005	32
16	Robert Hill	F	05/08/1982	RE	18/08/2008	26
17	Caleb Grant	F	28/02/1971	RE	28/10/2008	37
18	Shawn McCarty	F	28/02/1956	RE	31/05/2006	50
19	Jennifer Callahan	M	13/11/1975	RE	12/04/2006	30
20	Alexis Cortez	F	11/05/1954	LE	21/11/2008	54
21	Daniel Roberts	M	05/11/1954	RE	21/07/2006	51
22	Susan Howe	F	22/09/1954	RE	11/06/2008	53
23	Vanessa Moore	F	01/10/1952	RE	13/05/2010	57
24	Brittany Poole	M	15/10/1954	LE	31/07/2007	52
25	Angela Cunningham	F	06/06/1991	RE	06/06/2009	18
26	Joseph Robinson	M	07/10/1981	LE	02/10/2006	24
27	Jordan Reid	F	02/11/1983	LE	14/06/2005	21
28	Jordan Dunlap	M	01/04/1980	LE	05/02/2006	25
29	Dr. Susan Townsend	M	19/04/1984	RE	14/09/2006	22
30	Rebecca Malone	M	03/03/1997	RE	03/03/2015	17

Can you spot any problems in this RCT dataset?

<https://jamanetwork.com/journals/jamaophthalmology/fullarticle/28115>

	A	B	C	D	E	F
1	Surname and Name	Sex	Birthdate	Eye	Date of Surgery	Age at Sur
2	Alison Green	M	05/06/1976	RE	11/08/2008	32
3	Jared Friedman	F	13/04/1997	LE	13/04/2015	17
4	Jessica Cohen	M	21/03/1976	RE	28/12/2008	32
5	Sara Young DDS	F	12/05/1964	LE	03/04/2010	45
6	Whitney Douglas	M	26/01/1974	RE	02/01/2009	34
7	Daniel Watkins	M	13/11/1987	RE	03/11/2006	18
8	Jennifer Johnson	M	05/08/1952	RE	31/07/2010	57
9	Tara Thompson	M	16/06/1999	RE	16/06/2017	18
10	Jennifer Dixon	M	30/11/1977	RE	14/03/2007	29
11	Susan Howe	M	19/04/1955	LE	04/04/2005	49
12	Daniel McCarthy	F	11/10/1989	LE	30/04/2009	19
13	Bianca Boyd	M	07/11/1951	RE	02/05/2005	53
14	Lisa Kaufman	M	23/11/1996	RE	23/11/2014	17
15	Michael Stephen	M	06/10/1972	RE	21/07/2005	32
16	Robert Hill	F	05/08/1982	RE	18/08/2008	26
17	Caleb Grant	F	28/02/1971	RE	28/10/2008	37
18	Shawn Mccarty	F	28/02/1956	RE	31/05/2006	50
19	Jennifer Callahar	M	13/11/1975	RE	12/04/2006	30
20	Alexis Cortez	F	11/05/1954	LE	21/11/2008	54
21	Daniel Robertsor	M	05/11/1954	RE	21/07/2006	51
22	Susan Howe	F	22/09/1954	RE	11/06/2008	53
23	Vanessa Moore	F	01/10/1952	RE	13/05/2010	57
24	Brittany Poole	M	15/10/1954	LE	31/07/2007	52
25	Angela Cunningh	F	06/06/1991	RE	06/06/2009	18
26	Joseph Robinson	M	07/10/1981	LE	02/10/2006	24
27	Jordan Reid	F	02/11/1983	LE	14/06/2005	21
28	Jordan Dunlap	M	01/04/1980	LE	05/02/2006	25
29	Dr. Susan Townse	M	19/04/1984	RE	14/09/2006	22
30	Rebecca Malone	M	03/03/1997	RE	03/03/2015	17


Can you spot any problems in this RCT dataset?

- No correspondence between name and sex at all (e.g. Alison, Jessica, Whitney, Jennifer, Bianca all Male).
- This dataset was actually created using artificial intelligence (Chat GPT)
- Multiple eyes per person (fine), but we have people with two right eyes, people with different birthdays for each eye, and one person with five eyes
- (see M Wjst <https://tinyurl.com/3dmzfz8r>)

<https://jamanetwork.com/journals/jamaophthalmology/fullarticle/281150>


Example – dubious data

2-arm RCT




Example – dubious data

2-arm RCT




Example – dubious data

2-arm RCT




Other problems can be revealed by these plots



Take a moment – can you spot any problems?

Other problems can be revealed by these plots




Correlation across rows in the dataset

- We don't expect to see substantial correlation between the baseline values of successive participants.
- E.g. each participant's duration of infertility shouldn't be related to the duration of infertility of the person recruited after them, or to the next person's, or the next person's etc.
- We do expect correlation across rows if someone has typed (fabricated) values into the column – people are poor random number generators.
- Certainly don't expect this to differ between randomised groups.


Study ID	InfertilityDuration
1	12
2	8
3	8
4	3
5	1
6	3
7	9
8	3
9	7
10	6
11	5
12	1
13	4
14	6
15	2

Autocorrelation plot

Treatment group




Control group




- Plot of correlation between duration of infertility values 1 row apart (Lag = 1), 2 rows apart (Lag = 2), 3 rows apart etc.

Autocorrelation plot

Treatment group




Control group




- Plot of correlation between duration of infertility values 1 row apart (Lag = 1), 2 rows apart (Lag = 2), 3 rows apart etc.
- In treatment group, there is a correlation between successive rows, which decays as we get further apart.

Autocorrelation plot

Treatment group




Control group




- Plot of correlation between duration of infertility values 1 row apart (Lag = 1), 2 rows apart (Lag = 2), 3 rows apart etc.
- In treatment group, there is a correlation between successive rows, which decays as we get further apart.
- Control group looks like genuine data – no serial correlation between rows.

Autocorrelation plot

Treatment group




Control group




- Plot of correlation between duration of infertility values 1 row apart (Lag = 1), 2 rows apart (Lag = 2), 3 rows apart etc.
- In treatment group, there is a correlation between successive rows, which decays as we get further apart.
- Control group looks like genuine data – no serial correlation between rows.
- The treatment group correlation is suspicious, and the difference between groups is more so.


Relationships between variables

- Are expected relationships between variables present?
- Hard to fake (unless you know what you are doing). LLMs don't do a great job.
- Requires contextual knowledge (e.g. should we expect relationship between gestational age and birthweight in a particular trial).
- Don't expect multivariate distribution to differ between randomised groups.





- An example where relationships are too perfect – study of a diet intervention
- Data shown are pre-randomisation dietary variables



- An example where relationships are too perfect – study of a diet intervention
- Data shown are pre-randomisation dietary variables
- E.g. carbohydrate consumption perfectly correlated with fat consumption
- Implies that all participants consume carbohydrate and fat in the same ratio
- Statistical methods such as Principal Components Analysis can be useful in some cases (relationships between variables are too perfect)

The value of raw data

- These could be **easily detected** as being problematic with the IPD
- Suggests journals should request data for review with submissions
- Carlisle – increase in RCTs ‘critically flawed by false data’ when IPD inspected
- Better yet: publication of data with study – detection, deterrence
- How? INSPECT-IPD (Calvin Heal, NIHR Doctoral Research Fellow)

Original Article

False individual patient data and zombie randomised controlled trials submitted to *Anaesthesia*

J. B. Carlisle^{1,2} 

THE PREVALENCE OF ‘ZOMBIE’ TRIALS

More than one-quarter of a subset of manuscripts describing randomized clinical trials submitted to the journal *Anaesthesia* between 2017 and 2020 seemed to be faked or fatally flawed when their raw data could be examined, editor John Carlisle reported. He called these ‘zombies’. But when their raw data could not be obtained, Carlisle could label only 1% as zombies.

■ OK ■ Flawed data ■ Zombie

Raw data examined



Raw data not available



The value of raw data

- These could be **easily detected** as being problematic with the IPD
- Suggests journals should request data for review with submissions
- Carlisle – increase in RCTs ‘critically flawed by false data’ when IPD inspected
- Better yet: publication of data with study – detection, deterrence
- How? INSPECT-IPD (Calvin Heal, NIHR Doctoral Research Fellow)



INSPECT-AI

INveStigating ProBlEmatic Clinical Trials with AI

Start investigating, Goutham

Upload clinical trial PDFs to run automated integrity checks against INSPECT-SR tool.

 Start INSPECT-AI

 Open INSPECT-SR Tool

Avenell and colleagues, University of Aberdeen,
Vorland, Indiana University

ORIGINAL ARTICLE

Closed-loop automated oxygen control in late preterm and term, ventilated infants: A randomised controlled trial

Ourania Kaltsoyianni^{1,2} | Allan Jenkinson^{1,2} | Christopher Harris^{1,2} | Eleanor Jeffreys² | Oishi Sikdar² | Anne Greenough¹ | Theodore Dassios^{1,2}

¹Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, UK
²Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, London, UK

Correspondence
Theodore Dassios, Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, Denmark Hill, London SE5 9RS, UK.
Email: theodore.dassios@kcl.ac.uk

Funding information
National Institute for Health Research (NIHR) Biomedical Research Centre; Guy's and St Thomas' NHS Foundation Trust; King's College London

Abstract

Aim: To compare the time spent above the target oxygen saturation range ($SpO_2 > 96\%$) and the duration of supplemental oxygen between ventilated infants receiving closed-loop automated oxygen control (CLAC) or manual oxygen control in late preterm and term ventilated infants.

Methods: Infants were randomised to receive CLAC or manual oxygen control from recruitment and within 24 h of mechanical ventilation until successful extubation.

Results: Forty infants with a median (IQR) gestational age of 37.4 (35.9–38.5) weeks were studied at a corrected postmenstrual age of 37.6 (36.0–38.7) weeks. In infants randomised to CLAC ($n = 18$) the time spent above the target oxygen saturation range was reduced by 20% ($p < 0.001$), and the time spent in the target range (92%–96%) was increased by 32% ($p < 0.001$) and the time spent in hyperoxia was reduced ($p = 0.003$). CLAC reduced the time spent in hypoxemia ($SpO_2 < 85\%$) ($p = 0.017$) and there were fewer manual adjustments to the inspired oxygen concentration (FiO_2) ($p < 0.001$). There was no significant difference in the duration of supplemental oxygen ($p = 0.271$).

Conclusion: CLAC in ventilated infants born at or near term was associated with reduced time spent in hyperoxemia, more time spent in the target oxygen range, and fewer manual adjustments to the FiO_2 .



Does the study have an associated expression of concern or other relevant post-publication notice?

0

EOC/Correction Notices

0

PubPeer Comments

Expression of Concern & Correction Notices

No Concerns Found

SEARCH METHOD
DOI-based lookup

DOI (USED FOR SEARCH)
10.1111/apa.17549

TITLE
Closed-loop automated oxygen control in late preterm and term, ventilated infants: A randomised controlled trial

PubPeer Community Comments

No PubPeer Comments Found

This publication has no comments on PubPeer

Does the study have an associated expression of concern or other relevant post publication notice?

INSPECT-AI Suggestion: No

YesNoUnclearN/A

Comments

No comments on webpage either

Received: 27 September 2024 | Revised: 6 December 2024 | Accepted: 9 December 2024

DOI: 10.1111/apa.17549

ORIGINAL ARTICLE



Closed-loop automated oxygen control in late preterm and term, ventilated infants: A randomised controlled trial

Ourania Kaltsoyianni^{1,2} | Allan Jenkinson^{1,2} | Christopher Harris^{1,2} | Eleanor Jeffreys² | Oishi Sikdar² | Anne Greenough¹ | Theodore Dassios^{1,2}

¹Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, UK

²Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, London, UK

Correspondence
Theodore Dassios, Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, Denmark Hill, London SE5 9RS, UK.
Email: theodore.dassios@kcl.ac.uk

Funding information
National Institute for Health Research (NIHR) Biomedical Research Centre; Guy's and St Thomas' NHS Foundation Trust; King's College London

Abstract

Aim: To compare the time spent above the target oxygen saturation range ($SpO_2 > 96\%$) and the duration of supplemental oxygen between ventilated infants receiving closed-loop automated oxygen control (CLAC) or manual oxygen control in late preterm and term ventilated infants.

Methods: Infants were randomised to receive CLAC or manual oxygen control from recruitment and within 24 h of mechanical ventilation until successful extubation.

Results: Forty infants with a median (IQR) gestational age of 37.4 (35.9–38.5) weeks were studied at a corrected postmenstrual age of 37.6 (36.0–38.7) weeks. In infants randomised to CLAC ($n=18$) the time spent above the target oxygen saturation range was reduced by 20% ($p < 0.001$), and the time spent in the target range (92%–96%) was increased by 32% ($p < 0.001$) and the time spent in hyperoxia was reduced ($p=0.003$). CLAC reduced the time spent in hypoxemia ($SpO_2 < 85\%$) ($p=0.017$) and there were fewer manual adjustments to the inspired oxygen concentration (FiO_2) ($p < 0.001$). There was no significant difference in the duration of supplemental oxygen ($p=0.271$).

Conclusion: CLAC in ventilated infants born at or near term was associated with reduced time spent in hyperoxemia, more time spent in the target oxygen range, and

Q1.3 Research team concerns

Ctrl + /

Do other studies by the research team highlight causes for concern (retraction, expression of concern, relevant post-publication notices)?

7

Authors with No Mentions

0

Authors with Some Mentions

> Authors without Mentions in Retraction Watch (7)

Do other studies by the research team highlight causes for concern (retraction, expression of concern, relevant post-publication notices)?

INSPECT-AI Suggestion: No

Yes

No

Unclear


N/A

Clear

Comments

No concerns

Save comment



“Wait a minute! Isn’t anyone here a real sheep?”

Source: The Far Side, Gary Larson

What’s next?

- INSPECT-SR has been designed for RCTs, post-publication review, assuming no raw data access
- If you have raw data: INSPECT-IPD in development (Calvin Heal)
- Adapting for journal context: INSPECT-JR (Zewen Lu)
- Other study designs (e.g. Nonrandomised studies - INSPECT-NRS)
- Training and continual updating of guidance (new software, new research)

Trustworthiness assessment: everybody's doing it



Created with Google Gemini

- General awareness of research misconduct is probably **higher than ever**
- **This is good** – we want people to routinely consider trustworthiness of evidence
- But this is **new territory** for most people
- And while it isn't too hard to do well, it is **very easy to do it poorly** – so we need to use and promote good methods in our practice and guidance
- **Some** methods endorsed by **some** sleuths are just wrong – (this is not to denigrate the good work of the sleuthing community!) – statistical expertise not well represented
- Let's try to spread awareness of the issue, and also good methods for forensic analysis

Thanks, questions?